INNOVAX®-ND-ILT

Infectious Laryngotracheitis – Marek's Disease – Newcastle Disease Vaccine

Product Description

Safe, effective protection in a single vaccine against three of the most infectious and commercially disruptive flock diseases – infectious laryngotracheitis, Newcastle disease, and Marek's disease.

Indications

This product has been shown to be effective for the vaccination of healthy 18 day-old chicken embryos or one day-old chickens against Marek's disease, Newcastle disease and infectious laryngotracheitis.

-

Vaccination Programs

Many factors must be considered in determining the vaccination program for a particular farm or poultry operation. To be fully effective, the vaccine must be administered to healthy receptive birds held in proper environment under good management. In addition, the response may be modified by the age of the birds and their immune status. Seldom does 1 vaccination under field conditions produce complete protection for all individuals in a given flock. The amount of protection required will vary with the type of operation and the degree of exposure that a flock is likely to encounter.

Administration

In ovo Administration:

1. Inoculate each 18-day-old chicken embryo with a full dose (0.05 ml or 0.1ml).
2. Entire contents of container must be used within 1 hour after mixing or be discarded according to caution statement No. 11.
3. After reconstitution, the vaccine should be kept cool and gently agitated frequently.

Subcutaneous Administration:

1. Hold the chicken by the back of the neck just below the head. The loose skin in the area is raised by gently pinching with the thumb and forefinger. Insert the needle beneath the skin in a downward direction away from the head. Inject 0.2 ml per chicken.
2. Avoid hitting the muscles and bones in the neck.

3. Entire contents of container must be used within 1 hour after mixing or be discarded according to caution statement No. 11.

Supplied

1 x 2000 dose ampules

Caution

Good management practices are recommended to reduce exposure to Marek's disease and Infectious Laryngotracheitis for at least three weeks following vaccination. Therefore, directions should be followed carefully.

1. Do not mix any substance with this vaccine. Do not mix with other products, except as specified on this label.
2. Store vaccine in liquid nitrogen at a temperature below -150°C (-238°F).
3. Gloves and visor should be worn when handling liquid nitrogen.
4. **ONCE THAWED, THE PRODUCT SHOULD NOT BE REFROZEN.**
5. Do not dilute or otherwise stretch the dosage of this vaccine.
6. Once mixed with diluent, the vaccine should be gently agitated frequently.
7. Once mixed with diluent, the vaccine should be used within 1 hour.
8. Only healthy chicken embryos should be vaccinated.
9. Do not vaccinate within 21 days before slaughter.
10. This vaccine contains Gentamicin as a preservative.
11. Inactivate unused contents before disposal.
12. In case of human exposure, contact a physician.
13. **FOR ANIMAL USE ONLY.**