



Bivalent Avian Vaccine, Influenza Inactivated(H5N1 Re-5+H9N2 Re-2)

[Approval No] SYSZ (2006)150132125



MAC

250ml

Used only under the guidance of a veterinarian

■ Production Date:
■ Batch Number:
■ Expiration Date:

■ [Indication]: It is used to prevent Avian Influenza (H5 and H9).
■ [Method and Usage]: Subcutaneous injection in neck or muscular injection in chest with 0.3ml/ Bird for chicken of 2-5weeks old and 0.5 ml/bird for chicken over 5 weeks old.
■ [Storage & Validity]: 2-8 °C for one year

Produced by Qingdao Yebio Bioengineering Co.,Ltd , China

B I O E N G I N E E R I N G



For Veterinary Use Only

Instruction of Bivalent Avian Influenza Vaccine , Inactivated (H5N1 Re-5+ H9N2 Re-2)

[Approval No] SYSZ (2006) 150132125

[Product Name]

Common name: Inactivated Bivalent Avian Influenza (H5+H9)Vaccine(H5N1 Re-5+ H9N2 Re-2)

Commodity name: None

English name: Bivalent Avian Influenza(H5+H9) Vaccine, Inactivated (H5N1Re-5+ H9N2 Re-2)

Chinese name: Qinliugan (H5+H9) Erjia Michuoyimiao (H5N1Re-5+ H9N2 Re-2 zhu)

[Composition]

The vaccine is composed of inactivated reassortant avian influenza virus (H5N1Re-5+ H9N2 Re-2):(H₂N₂). The HA titer of chicken embryo fluid before inactivation is $\geq 9 \log_2$.

[Properties] Milky white emulsion.

[Actions and Uses]

It is used to prevent avian influenza virus (H5 Subtype) infection.

[Application and Dosages]

Subcutaneous injection in neck or intramuscular injection in chest ,0.3 ml/bird for 2~5-Week-old chickens, 0.5 ml/bird, for plus chickens above 5-week-old .

[Abnormal Reaction] There are no abnormal reactions generally.

[Precautions]

- (1) The product is not allowed to use in AIV infected chickens or in chickens with abnormal health conditions.
- (2) The product can never be frozen.
- (3) The product can not be used when there are damages, foreign materials or emulsion breakage and stratification.
- (4) The vaccine should be recovered to normal temperature and evenly mixed by shaking before use.
- (5) Sterile equipment should be used when vaccination Needles should be changed time -ly. It is advisable to have one needle for one chicken.
- (6) The vaccine should be finished in the day when unbottled.
- (7) It can't be used for the chickens within 28 days before slaughter.

[Specification] 100ml/bottle,250 ml/bottle, 500 ml/bottle.

[Package] 100bottle/box,40 bottles/box, 20 bottles/box.

[Storage&Expiration] The product can be stored at 2~8 °C for 12 months.

Used as instructed by a veterinarian

Qingdao YEBIO Bioengineering Co.,Ltd

Add: 25 Shanghai Road Huangdao, Qingdao 266500 P.R.china

Sales: 0086-532-85663027 85620449

Fax:0086-532-85663027

E-mail: YEBIO@YEBIO.com.cn Website:http://www.YEBIO.com.cn

YEBIO Bioengineering co.,ltd of QINGDAO