



MAC

NYBH[2007]47

Reassortant Bivalent Avian Influenza Virus Vaccine,
Inactivated (H5N1 Subtype, Re-5+Re-4 Strain)

250ml



■ Production Date:

■ Batch Number:

■ Expiration Date:

■ Indication: It is used to prevent avian influenza caused by H5 subtype avian influenza virus.

■ Method and Usage: Subcutaneous injection in neck or muscular injection in chest, 0.3ml/bird for chickens between 2-5 weeks old; 0.5ml/bird for chickens above 5 weeks old.

■ Storing Condition: 2-8 °C for one year.

Used Only under the Guidance of a Veterinarian

Produced by Yebio Bioengineering Co., Ltd of Qingdao, China

B I O E N G I N E E R I N G



For Veterinary Use Only

**Reassortant Bivalent Avian Influenza Virus Vaccine,
Inactivated (H5N1 Subtype, Re-5+Re-4 Strain)**

[Approval No] NYBH [2007] 47

[Product Name]Common name: Reassortant Bivalent Avian Influenza Virus Vaccine, Inactivated
(H5N1 Subtype, Re-5+Re-4 Strain)

Commodity name: None

English name: Reassortant Bivalent Avian Influenza Virus Vaccine, Inactivated
(H5N1 Subtype, Re-5+Re-4 Strain)

Chinese name: Qinliuganbingdu Erjia Miehuoyimiao (H5N1 Yaxing, Re-5 zhu+Re-4 zhu)

[Composition]The vaccine is composed of inactivated reassortant avian influenza virus H5N1 Subtype, Re-5 and Re-4 strain. The HA titer of the chicken embryo fluid of Re-5 strain before inactivation is $\geq 9\log_2$, and the HA titer of the chicken embryo fluid of Re-4 strain is $\geq \log_2$.**[Property]** Milky white emulsion.**[Actions and Uses]**

It is used to prevent avian influenza due to (H5 Subtype) AIV.

[Application and Dosages]

Subcutaneous injection in neck or intramuscular injection in chest, 0.3 ml/bird for 2~5 Weeks old chickens, 0.5 ml/bird for over 5 weeks old chickens.

[Abnormal Reaction] Generally no abnormal reactions.**[Precautions]**

- (1) The product cannot be used for chickens infected by AIV or with abnormal health condition.
- (2) Freezing is prohibited.
- (3) The product can not be used if packing is damaged, impurity is found or emulsion is Broken or stratified.
- (4) Make sure the vaccine is at room temperature and is well shaken before use.
- (5) Sterile equipment should be used in vaccination. Needles should be changed timely. It is suggested that a separate needle be used for each injection.
- (6) The vaccine should be finished the same day when unbottled.
- (7) It can't be used for the chickens within 28 days before slaughter.

[Specification] 100ml/bottle, 250 ml/bottle, 500 ml/bottle.**[Package]** 100bottles/box, 40 bottles/box, 20 bottles/box.**[Storage&Expiration]** 2~8°C for one year.

Used as instructed by a veterinarian

YEBIO Bioengineering Co., Ltd of Qingdao

Add: 25 Shanghai Road Huangdao, Qingdao 266500 P.R.china

Sales: 0086-532-85663027 85620449

Fax: 0086-532-85663027

E-mail: YEBIO@YEBIO.com.cn Website: <http://www.YEBIO.com.cn>

YEBIO Bioengineering co., ltd of QINGDAO