

AVIPRO® 304 ND-IB-MG

L.A.H.I.

NEWCASTLE-BRONCHITIS VACCINE

MASS. TYPE, KILLED VIRUS

MYCOPLASMA GALLISEPTICUM BACTERIN

FOR VETERINARY USE ONLY

Directions For Use: Vaccine should be warmed to room temperature before use. Shake well before prior to and during use.

Inject by subcutaneous inoculation in the neck using aseptic technique. Use entire contents of bottle when first opened.

Administered to birds at 6 weeks of age or older. At a minimum of 3 weeks prior to the administration of the first dose of killed vaccine, a live Newcastle-Bronchitis vaccine (Mass. type) must be used for priming.

Revaccination with the killed vaccine should be done 3-6 weeks after the first vaccination. Vaccinate only healthy birds.

Dose: 1/2 ml (0.5 ml) per bird. Inject all birds in flock.

Vaccine was carefully produced and passed all tests in accordance with the U.S. Government.

Recommended for the vaccination of chickens as an aid in the prevention of infectious bronchitis (Mass. type) and Newcastle disease. It is also used as an aid for the control of clinical signs associated with M. gallisepticum infection in chickens.

Caution: Store at 2°-7°C (35°-45°F). Do not freeze.

Warning: In case of accidental human injection seek immediate medical attention.

Do not vaccinate within 42 days before slaughter.

Manufactured in the USA by: **LOHMANN ANIMAL HEALTH INT.**, Winslow, Maine 04901 U.S.A.

U.S. Vet. Lic. No. 196

	LIST NO.
1000 doses	1831

NAC No.: 10081181