



MAC

250ml Infectious Coryza and Newcastle Disease Vaccine, Inactivated

SYSZ (2006) 150132044



制涂层 辨真伪

请仔细阅读说明
SHEJIAOYU LIJIAOYU
SHEJIAOYU LIJIAOYU
SHEJIAOYU LIJIAOYU

- Production Date:
- Batch Number:
- Expiration Date:

- **Indication:** The product is prepared with imprinted adjuvant, and is used to prevent Infectious Coryza and ND.
- **Method And Usage:** Chickens are subcutaneously injection in neck: 0.25ml/bird for chickens between 24 and 42 days old low 2 weeks old; 0.5ml/bird for chickens over 2 months old;
- **Storage & Validity:** 2-8 °C for one year.

Qingdao Yebio Bioengineering Co., Ltd of , China

B I O E N G I N E E R I N G



For Veterinarian Use Only

Instruction Infectious Coryza and Newcastle Disease Vaccine, Inactivated

Approval No: SYSZ (2006) 150132044

[Product Name]

Generic Name: Infectious Coryza and Newcastle Disease Vaccine, Inactivated

Trade Name: None

English Name: Infectious Coryza and Newcastle Disease Vaccine, Inactivated

Chinese Alphabetic: Ji Chuanranxingbian; Ji Xinchengyi ErlianMiehuoyimiao

[Composition] The product is prepared with imported adjuvant. The vaccine contains inactivated Hemophilus paragall: A C-Hpg-8, C-Hpg-668 and NDV LaSota strain. Before inactivation the ND virus concentration is $\geq 10^8$ EID₅₀/0.1ml; Hemophilus paragalli A and C are $\geq 24 \times 10^8$ CFU/ml.

[Property] Cream white emulsion

[Indication] It is used to prevent Infectious Coryza and ND. Immunity appears 14-21 days after vaccination. One vaccination has an immune duration of 3-5 months; first vaccination at 21 days old followed by a second one at 120 days old has an immune duration of a months.

[Application & Dosage] Chickens are subcutaneously injected in the neck 0.2ml/bird for chickens between 24 and 42 days old; 0.5ml/bird for chickens above 42 days old.

[Abnormal reaction] no

[Precaution]

- (1) Freezing is prohibited, and the frozen vaccine is not allowed to use.
- (2) The vaccine should be recovered to normal temperature and mixed evenly prior to use.
- (3) The vaccine should be finished in the same day of unbottling..
- (4) The product is exclusively used in the health poultry.
- (5) The injection site should be disinfected locally.
- (6) The contaminated items such as used bottles, remaining vaccines, utensils should be disposed by disinfection treatment.
- (7) The product is not allowed to use in meet poultry within 21 days before slaughter, and in other poultry within 42 days before slaughter.

[Specification] 100ml/bottle, 250ml/bottle, 500m/bottle.

[Package] 100bottles/box, 40bottles/box, 20bottles/box.

[Storage & Expiration] 2-8°C, one year.

Used as instructed by a veterinarian

Qingdao YEBIO Bioengineering Co., Ltd

Add: 25 Shanghai Road Huangdao, Qingdao 266500 P.R. china

Sales: 0086-532-85663027 85620449

Fax: 0086-532-85663027

E-mail: YEBIO@YEBIO.com.cn Website: http://www.YEBIO.com.cn

YEBIO Bioengineering co., ltd of QINGDAO